

NIMUE MD RANGE

EDUCATION ONE PAGER

SCAR GEL | 30ML

A gel for the prevention of scarring post-surgery and trauma, assisting in the reduction of most types of existing scars. It also assists in reducing PIHP. The inclusion of all the active ingredients in the right concentrations is very effective in the prevention and reduction of scar tissue formation. Commencing treatment within 2 weeks post surgery or trauma is recommended.

INDICATIONS

- Scars (including post-surgical scars and acne scars)
- Trauma
- Burns
- Cuts
- Tattoo removal

BENEFITS

- Inhibits stress related inflammation and reduces skin redness
- Stimulates the production of collagen
- Repairs and accelerates wound healing
- Improves tissue microcirculation
- Assists in reducing and preventing PIHP

KEY INGREDIENTS

- Epidermal Growth Factor (sh-Oligopeptide-1) (*Accelerates healing and cell renewal*)
- Turmeric Plant Stem Cells (*Neutralises inflammation and accelerates wound healing*)
- Lipohydroxy Acid (*Gentle exfoliation and skin renewal*)
- Anti-Stretch Complex (*Stimulates cell growth, tissue repair and collagen synthesis*)
- Glycyrrhetic Acid (*Controls redness and soothes a sensitised skin*)

DIRECTIONS FOR USE

SCAR GEL	
(Use following suture removal or when skin has closed post injury or trauma)	
Homecare use	
Face, neck and décolleté	
2 Pumps minimum	
Usage guideline	
Area	Pumps
10cm scar length	1 pump
<ul style="list-style-type: none">• Apply surgical tape to the area and apply MD Scar Gel to the surface of the tape 2-3 times daily, massaging gently• Wipe tape clean before re-application of MD Scar Gel• Replace tape every 4-5 days• Use on its own and do not mix with other products• DO NOT USE ON EZCEMA, PSORIASIS OR SKIN CANCER	



MD

NIMUE MD RANGE

BEFORES AND AFTERS

SCAR GEL | 30ML

BEFORE



WEEK 2



BEFORE



WEEK 6



Case studies performed in clinical practice by a Plastic Surgeon
For further case studies, please refer to MD information booklet

MD